Thrusts and Prospects on Understanding and Predicting Asian Monsoon Climate*

WANG Bin†(王 斌)

Department of Meteorology and International Pacific Research Center, School of Ocean and Earth Science and Technology, University of Hawaii at Manoa, Honolulu, Hawaii 96822, *USA*

(Received October 7, 2008)

ABSTRACT

Development of monsoon climate prediction through integrated research efforts to improve our understanding of monsoon variability and predictability is a primary goal of the Asian Monsoon Years (2007–2011) and International Monsoon Study under the leadership of the World Climate Research Programme. The present paper reviews recent progress in Asian monsoon research focusing on (1) understanding and modeling of the monsoon variability, (2) determining the sources and limits of predictability, and (3) assessing the current status of climate prediction, with emphasis on the weekly to interannual time scales. Particular attention is paid to identify scientific issues and thrust areas, as well as potential directions to move forward in an attempt to stimulate future research to advance our understanding of monsoon climate dynamics and improve our capability to forecast Asian monsoon climate variation.

Key words: Asian monsoon, climate predictability, climate prediction, intraseasonal oscillation

1. Introduction

The giant Asian monsoon system embodies the most complex interactions between Earth's atmosphere, hydrosphere, cryosphere, and biosphere including human activities. The Asian monsoon interacts with the El Niño-Southern Oscillation (ENSO) and underlying oceans, resulting in far-reaching impacts on global climate and environment. Given the necessity of reducing uncertainty in global warming projections, understanding the dynamics of monsoon systems is of paramount importance.

The Asian monsoon variability significantly influences the economy and society across Asian countries, where about 50% of the world population inhabits. Any improved knowledge about monsoon rainfall variation and future change will be of great importance to sustainable development of world economy.

Monsoon science has advanced enormously in the last two decades. These advances have been comprehensively reviewed by leading scientists in the recent publications (e.g., Webster et al., 1998; Chang et al., 2005; Wang, 2006). Unprecedented amount of data derived from satellite observations and field experiments and the advance in improvement of climate models, computer power, and communication technology have deepened our understanding of the monsoon phenomena and enhanced operational capability in monsoon prediction.

Despite significant societal and environmental demands for accurate monsoon prediction and the notable improvements in our ability to simulate the Asian monsoon, operational prediction of Asian monsoon variations is still in its infancy and its achievement is seen as a great challenge faced by operational weather forecast centers worldwide.

It is imperative to improve Asian monsoon prediction through integrated research efforts to enhance our understanding of Asian monsoon variability and predictability. This is the main goal of the Asian Monsoon Years (AMY 2007–2011), which is a cross-cutting initiative as part of the International Monsoon Study (IMS) under the leadership of the World Climate Research Programme (WCRP).

This article reviews progresses in Asian monsoon research focusing on our capability in numerical

*Supported by NSF climate dynamics program (ATM06-47995) and by IPRC, which is in part sponsored by FRCGC/ JAMSTEC, NASA, and NOAA.

†Corresponding author: wangbin@hawaii.edu.

modeling of climate variability and prediction of the prominent monsoon active/break periods and year-to-year fluctuations. Since increased knowledge in the Asian monsoon variability facilitates improvement of our modeling and predicting capability, the current understanding of the Asian monsoon variability will also be reviewed. Particular attention is paid to discuss major scientific issues and road blocks in an attempt to stimulate future research.

Section 2 discusses the monsoon responses to external solar forcing: the diurnal and annual cycles. Sections 3 and 4 deal with Asian monsoon variability, predictability, and prediction on the intraseasonal and interannual time scales, respectively. Different from the forced responses, the intraseasonal-interannual variations are due to internal feedback processes within the atmosphere and/or the coupled climate system. The last section briefly discusses some emerging modeling and prediction issues and future directions.

This review will emphasize precipitation—the most important variable for gauging the monsoon variability and making monsoon prediction. Precipitation is also one of the measures of the global water and energy cycle and holds a key in linking external radiative forcing and the atmospheric general circulation. Efforts are also made to merge the discussions of the knowledge and understanding with the discussions of modeling, predictability and prediction.

2. Monsoon diurnal and annual cycles

Understanding and modeling the diurnal and annual cycles are essentially important for a number of reasons. Both the annual and diurnal cycles play significant roles in regulating monsoon variations in rainfall and circulation, thus affecting weather and climate fluctuations. The performance of numerical models in simulating seasonal mean states is closely related to their capability to predict seasonal anomalies (Sperber and Palmer, 1996; Slingo et al., 1996). Getting the annual cycle right is of critical importance for models to reproduce accurate teleconnection and climate anomalies away from the ENSO region (Bengtsson et al., 1993). The diurnal and annual cycles are also most relevant for revealing monsoon modeling problems because both of them have the largest amplitudes in the monsoon regions. Moreover, the diurnal cycles provide an efficient way for verification of models' physical parameterizations (Yang and Slingo, 2001). In practice, the onset and withdrawal of summer monsoons are of vital importance as agriculture and other human activities are strongly influenced by the period and/or duration of monsoon rainy season.

2.1 Observed diurnal cycle of precipitation

Monsoon regions exhibit the largest magnitude and most complex behavior of the precipitation diurnal cycles (Kikuchi and Wang, 2008). It has been recognized for a long time that strong diurnal variation of rainfall is a basic feature in the tropical land regions. However, only since the satellite era begins, the global as well as minute diurnal features of precipitation can be revealed (e.g., Murakami, 1983; Nitta and Sekine, 1994). Comparisons of satellite and gauge data have been conducted in Indo-China in the GAME project (Ohsawa et al., 2001) and over China (Yu et al., 2007). Recently, accumulation of the Tropical Rainfall Measuring Mission (TRMM) satellite observations has enabled to capture microscopic features of the diurnal variations over the entire tropics (e.g., Sorooshan et al., 2002; Nesbitt and Zipser, 2003).

Using two complementary TRMM datasets (3B42 and 3G68) for the 1998–2006 period, Kikuchi and Wang (2008) revealed that, in addition to the oceanic diurnal regime with an early morning peak in rainfall and the continental regime with an afternoon peak, there is a universal coastal regime in global monsoon regions, which shows strong rainfall movements, either inland or offshore. Figure 1 presents the evolution of the costal pattern in South Asia and maritime continent. Over South Asia, the inland propagation sets in along the coastline of India and the Indo-China Peninsula at 0900 LST and then becomes stronger and moves inland around noon (1200–1500 LST), peaking around 1500 LST; finally, the rain stops farther inland around 2100 LST. In the coastal offshore propagation, rain tends to start near the coast around midnight, expand over the ocean to 1200 LST, and then disappear between 1200 and 1500 LST. The results shown in



Fig.1. Evolution of precipitation diurnal cycle reconstructed from the first two Extended EOF (EEOF) modes for (a) South Asia and (b) maritime continent. The corresponding modified (unmodified) local standard times are shown at the right corner of each subpanel. The fractional variances accounted by the two EEOF modes are shown in the titles. Modified from Kikuchi and Wang (2008).

Fig.1 provide a measure for gauging models' performance in simulation of the diurnal cycles.

Descriptions of the diurnal cycle using satellite data to date are primarily for clouds and rainfall; the related circulation and other fields have not been fully analyzed, except a few intensive observational analyses that provide detailed regional characteristics (e.g., Mori et al., 2004). We need to improve our understanding of the relationship between diurnal cycle and surface orography and land/sea configurations; the causes of the inland and offshore propagations of the diurnal cycle from coastal regions; the effects of the diurnal variation on the cloud/rainfall variations over the open ocean; and the interaction between the diurnal cycle and the intraseasonal oscillation (ISO).

2.2 Observed annual cycle of precipitation

Although conventional definition of the monsoon has been solely based on annual reversal of prevailing surface winds (Ramage, 1971), the monsoon climate is not only characterized by a change in surface winds, but also by its contrasting rainy summer and dry winter (Webster, 1987; Trenberth et al., 2000). The monsoon itself is a manifestation of the annual variation



Fig.2. (a) Normalized principal components of the first two multi-variable EOF modes of the climatological monthly mean precipitation and the winds at 850 hPa. (b) and (c) are their corresponding spatial patterns of precipitation (shading, unit: mm day^{-1}) and winds (vectors in units of m s^{-1}) at 850 hPa for EOF1 and EOF2, respectively. Winds with wind speed of less than 1 m s^{-1} are omitted. (d) Solstice mode of the annual variation as described by the differential precipitation rate (mm day^{-1}) and the 850 hPa winds (namely, JJAS minus DJFM). (e) Equinoctial asymmetric modes as described by the April-May mean minus the October-November mean precipitation rate (mm day^{-1}) and the 850 hPa winds. Winds with wind speed of less than 4 and 2 m s^{-1} are omitted in (d) and (e), respectively. Modified from Wang and Ding (2008).

of the tropical general circulation (e.g., Webster, 1987; Zeng and Li, 2002).

Wang and Ding (2008) has shown that the global monsoon can be quantitatively defined by the first two empirical orthogonal modes of annual variation in global precipitation and low-level (850 hPa) winds (Fig.2). Both modes have an annual period (Fig.2a). The first mode features an interhemispheric contrast in precipitation (Fig.2b) and can be simply described as a June-July-August-September minus December-January-February-March precipitation pattern (Fig. 2d). Thus, the first mode is called the solstice mode, which reflects the impact of antisymmetric annual solar forcing with a one-to-two-month phase delay in atmospheric response. The second mode has the maximum and minimum occurring around April and October, respectively. Its spatial pattern (Fig.2c) resembles the April-May (AM) minus October-November (ON) precipitation and circulation pattern (Fig.2e). Therefore, the second mode represents an equinoctial asymmetric mode, or spring-fall asymmetry, which is one of the important features regarding the seasonal variation in tropical and monsoon circulation. As shown in Fig.2, the primary features of the annual cycle of tropical circulation can be quantitatively defined by a combination of the solstice mode and the equinoctial asymmetric mode; together, they account for 84% of the annual variance.

Although depiction of the rainy seasons over the land and islands has attracted numerous studies using rain gauge observations (e.g., Rao, 1976; Tao and Chen, 1987; Ding, 1992; Tanaka, 1992; Matsumoto, 1997), for a long time our knowledge has fallen short about the rainy season over the Asian marginal seas (such as the Arabian Sea, Bay of Bengal, South China Sea, and East China Sea). The precise seasonal march of the rainfall over Asia and adjacent oceans has been revealed only when accumulation of satellite derived proxy and global rainfall data are available (e.g., Wang, 1994; Lau and Yang, 1997).

Figure 3 (adopted from Wang and LinHo, 2002) shows that the large-scale onset of Asian monsoon rainy season starts from the Andaman Sea north of the Sumatera toward the end of April. The grand onset of Asian monsoon is characterized by the rainfall surges over the South China Sea (SCS) in mid-May,



Fig.3. Dates of onset determined by relative climatological pentad mean (CPM) rainfall. The thick solid lines represent the Asian-Pacific summer monsoon domain defined by the maximum CPM occurring in boreal summer (May-September) and the annual range of the CPM exceeding the January mean rainfall rate by 5 mm day^{-1} (and the local January mean rainfall rate itself). The onset pentad is the first pentad when the CPM rainfall exceeds the January mean rainfall rate by 5 mm day^{-1}. The thick dashed lines denote discontinuities (merger of three or more contours). The arrows point to the directions of rain-belt propagation. The thin dashed line denotes the location of the subtropical monsoon front formation, which divides East Asian subtropical monsoon and tropical western North Pacific monsoon where their onset patterns are entirely different. Adopted from Wang and LinHo (2002).

which establishes a planetary-scale monsoon rain band extending from the Bay of Bengal to the subtropical western North Pacific (WNP). The rain band then advances northwestward, initiating the continental Indian rainy season, the Chinese Meiyu, and the Japanese Baiu in early to mid-June.

It is not well understood why Asian monsoon first starts from the Andaman Sea and whether the monsoon onset is a manifestation of instability of the changing mean flow, or, it is caused by a finite amplitude disturbance that is formed after large scale mean flow is pre-conditioned. On the other hand, pre-monsoon rainfalls and latent heat release are seen in southern China (Tao and Chen, 1987; Ding, 2004), Indo-China (Kiguchi and Matsumoto, 2005) and over the Tibetan Plateau (Ueda et al., 2003; Taniguchi and Koike, 2007). The upper tropospheric heating is enhanced by such pre-monsoon rainfall events. How important these pre-monsoon rainfalls are and how they affect interannual variation of the monsoon onset are still open questions.

Although solar radiation forcing has sinusoidal variation, the seasonal cycle of the Asian monsoon often experiences stepwise, abrupt changes (e.g., Matsumoto, 1992; Ding, 1994, 2004; Ueda and Yasunari, 1996). The monsoon onset often occurs abruptly. For instance, Fig.3 shows a three-stage stepwise onset over the WNP. Wang and LinHo (2002) have shown that the heights of the rainy seasons occur primarily in three stepwise phases: in late June over the Meiyu/Baiu regions and the northern Bay of Bengal; in late July over India and northern China; and in mid-August over the tropical WNP. Another important aspect of the monsoon seasonal cycle is the asymmetric nature of the seasonal transition between boreal spring and autumn (Matsumoto and Murakami, 2002; Hung et al., 2004; Chang et al., 2005). In general, there is a need to identify the principal physical processes that determine the abrupt evolution in seasonal cycle, although the nonlinear atmospheric processes and the atmosphere-ocean interactions (Wu and Wang, 2001; Ueda, 2005) and land-atmosphere-ocean interaction processes (Minoura et al., 2003) have been hypothesized as the possible agents.

2.3 Numerical modeling of the diurnal and annual cycles

Accurate simulation and prediction of the spatial and temporal variation of diurnal rainfall around the globe remains one of the unsolved problems in climate system modeling. Sperber and Yasunari (2006) summarized the problems in the simulation by GCMs of the diurnal cycle. Adequate resolution of the planetary boundary layer, the coupling between the planetary boundary layer and deep convection may hold a key to the improvement of the diurnal cycle in GCMs.

What are the major weaknesses of the climate models in simulation of the annual cycle? Kang et al. (2002) found that all of 11 GCMs, which participated in the CLIVAR/Asian-Australian Monsoon Panel (AAMP) AGCM intercomparison project, overestimate the amplitudes of climatological seasonal variations of the Indian summer rainfall but underestimate the rainfall variation in the WNP. Recent evaluation of the 17 climate prediction models confirms that a common weakness lies in the WNP and East Asian monsoon region (Lee et al., 2008).

In the forced response of the monsoon, the following questions remain to be addressed: (a) What determines the structure and dynamics of the diurnal and annual cycles of the coupled atmosphere-ocean-land system? (b) How can the major weaknesses of climate models in simulation of the diurnal cycle and annual cycle of global precipitation be remedied?

3. Intraseasonal variability and predictability

The monsoon intraseasonal oscillation (MISO) is a dominant form of monsoon variability that links weather and climate. The MISO is closely related to the onset, active, and break periods of the monsoons. The Madden-Julian (1971, 1972) oscillation (MJO) has important impacts on MISO, but the behavior of the MISO is more complex than the MJO. This is in part due to the fundamental modification of the monsoon circulation on the MJO. In addition, the monsoon has its own intrinsic ISO modes (Krishnamurthy and Shukla, 2008). For instance, the 10–25-day modes and the independent northward

propagating mode: about one half of the northward propagating ISOs over the Indian Ocean are independent of the equatorial eastward propagating MJO (Wang and Rui, 1990; Hendon et al., 2007).

3.1 Origin of the monsoon intraseasonal oscillation

The characteristics of the observed MISO have been extensively reviewed by Waliser (2006). It has been generally recognized that the MISO has the following essential features: (a) northward propagation in the Indian monsoon region (Yasunari, 1979, 1980; Sikka and Gadgel, 1980) and northwestward propagation in the WNP (e.g., Lau and Chan, 1986; Nitta, 1987; Hsu and Weng, 2001); (b) formation of a northwest-southeastward tilted anomalous rain band near Sumatera (Maloney and Hartmann, 1998; Annamalai and Slingo, 2001; Kemball-Cook and Wang, 2001; Lawrence and Webster, 2002); (c) initiation in the western equatorial Indian Ocean (60°–70°E) (Wang et al., 2005a, 2006; Jiang and Li, 2005), (d) the phase-locking to the annual cycle, or climatological ISO (CISO) (Nakazawa, 1992; Wang and Xu, 1997; LinHo and Wang, 2002), and (e) the prominent 10–25-day oscillation in the off-equatorial South Asian monsoon trough (Krishnamurti and Bhalme, 1976; Chen and Chen, 1993; Wu and Zhang, 1998; Wen and Zhang, 2008). In addition, the MISO has close interaction with ISOs in the mid-latitude region (Kawamura et al., 1996) and extratropical wave trains (Ding and Wang, 2007) due to its proxy to the subtropics.

Theories have been proposed to explain the essential features of the MISO. A review is provided in Wang (2005). The northward propagation has been explained in terms of boundary layer destabilization-convective stabilization (Webster, 1983; Goswami and Shukla, 1984), air-sea interaction (Kemball-Cook and Wang, 2001; Fu et al., 2003), and the effects of the monsoon easterly vertical shear (Jiang et al., 2004; Drbohlav and Wang, 2005). The formation of the NW-SE tilted precipitation belt has been interpreted as resulting from the emanation of convectively coupled equatorial Rossby waves from the decaying equatorial MJO disturbances (Wang and Xie, 1997; Lawrence and Webster, 2002). The re-initiation of the monsoon active-break cycles was attributed to local SST and hydrological feedback (Stephens et al., 2004) and a self-induction mechanism (Wang et al., 2005a, 2006). The role of the topographic effect of the maritime continent was also discussed by Hsu et al. (2004) and Hsu and Lee (2005).

However, the MISO involves multi-scale interactions among diurnal cycle, mesoscale and synoptic scale disturbances, the MJO and annual cycle. Full interpretation of the MISO phenomena remains elusive. Many issues invite further investigations, for instance: (a) What are the typical multi-scale and 3D structures of the MISO? (b) Are multi-scale interactions essential for development and maintenance of the MJO and MISO and if so how? (c) Why is there a 10–25-day oscillation and how is it related to the MISO? (d) How do we get a complete theoretical framework for describing the characteristics of the MISO? (e) What roles do atmosphere-ocean-land interactions play in sustaining the MISO? (f) To what extent and how does the ISV in SST depend on atmospheric forcing? And how does intraseasonal SST anomaly feedback to the ISO?

3.2 Modeling of intraseasonal variations

The realistic representation of tropical convection in our global atmospheric models is a long-standing grand challenge for numerical weather forecasts and global climate predictions (Waliser, 2006). The capability of models in reproducing the MJO and MISO has been continuously assessed over the last decade. Pertaining to the MISO simulations, Sperber et al. (2001) found that the AGCMs have difficulty in representing the pattern of precipitation associated with the dominant mode; and also the models usually fail to project the subseasonal modes onto the seasonal mean anomalies. Waliser et al. (2003b) analyzed the MISO in the 10 AGCMs that participated CLIVAR/AAMP AGCM intercomparison project and found that (a) the most problematic feature is the overall lack of variability in the equatorial Indian Ocean, (b) most of the model ISO patterns did exhibit some form of northward propagation, but they often show a southwest-northeast tilt rather than the observed northwest-southeast tilt, and (c) the fidelity of a model to represent boreal summer versus winter intraseasonal variability (ISV) appears

to be strongly linked.

Why do AGCMs have considerable difficulties in simulation of the MJO and MISO? One of the major difficulties involves cumulus parameterization used to estimate the vertical redistribution of heat and moisture by unresolved convective clouds in GCMs (Slingo et al., 1996; Maloney and Hartman, 2001). In general, modeling of the ISO in the complex AGCMs must entail a series of parameterizations including moisture transport, clouds and convection, and radiation transfer; thus, uncertainties in mathematical descriptions of these multi-scale interactive parameterizations can jeopardize the model's capability in simulating the MJO (Wang, 2005). For instance, it is critical for models to get correct heating partitioning between convective and stratiform precipitation, yet AGCMs generally underestimate the portion of stratiform precipitation. The inadequate treatment of cumulus parameterization and the multi-scale interaction processes are the major hurdles for realistic simulation of MJO and MISO.

How to rectify systematic errors in models' simulation of the MJO remains a great challenge and many modeling issues remain unresolved. For instance, what are the roles of radiative heating and mesoscale systems in determining the heating profile (convective/stratiform) and how can this be correctly represented in models? How important is the modulation of the diurnal cycle in intraseasonal monsoon variations? How do the errors in simulating ISOs impact simulation of the interannual variability?

3.3 Predictability and prediction skills of MJO and MISO

The capability of daily weather forecast is limited by the chaotic nature of Earth's atmosphere and the growth of the initial errors (Lorenz, 1963) and useful weather forecast can be made only within a week or two. On the other hand, the statistical behavior of the weather averaged over a season may be predictable due to the persistent forcing from anomalous conditions at the sea surface and land surface (Charney and Shukla, 1981). The intraseasonal variability (ISV) falls between the daily weather and the seasonal mean climate. The ISV is largely governed by atmospheric internal dynamics (Palmer, 1994; Waliser et al., 2006) and is therefore, to a large extent, chaotic in nature and unpredictable. Fu et al. (2007) showed that air-sea coupling may extend ISV predictability, suggesting that atmosphere-ocean interaction may be a source of predictability for ISV. But in theory, the source and limiting factors for intraseasonal predictability remain elusive.

Based on the perfect model assumption, previous studies have suggested a theoretical forecast limit of ISO (potential predictability) out to 15 days for rainfall and to 30 days for upper level circulation field (e.g., Waliser et al., 2003b; Waliser, 2005; Liess et al., 2005; Fu et al., 2008; Pegion and Kirtman, 2008).

The estimated predictability and prediction skills of ISV were mostly made in terms of band-pass filtered MJO signals (a portion of ISV) or a portion of the MJO signals (the principal modes of the MJO). Using filtered data or principal modes for assess predictability/prediction skill are convenient but unconventional in comparison to assessing the weather and seasonal mean climate prediction. The true meaning of intraseasonal predictability and adequate metrics for assessing intraseasonal prediction remains to be rigorously defined.

About a decade ago, the dynamical forecast of the MJO by using the atmospheric only model of the vintage of the NCEP reanalysis had a useful skill only up to 9 days for boreal winter season (Hendon et al., 1999; Jones et al., 2000). However, dynamical models have improved significantly in the last decade (Sperber and Waliser, 2008) and in a few cases are able to predict the MJO out to a lead-time comparable to the empirical-statistical schemes (Seo et al., 2005; Kim et al., 2008; Vitart et al., 2008). Nevertheless, prediction of MJO is still in its infancy.

The prediction skills of the MISO have been assessed recently in the hindcast experiments made by DEMETER and APCC/CliPAS models (Kim et al., 2008). These models tend to considerably underestimate the intraseasonal variances over the WNP and Indian monsoon regions. Only a few models can realistically capture the evolution and structure of the

boreal summer ISO, such as the NW-SE slanted precipitation band. Although models have large systematic biases in the spatial pattern of dominant ISV, the leading EOF modes of the ISO in the models are closely linked to the models' ENSO, which is a feature that resembles the observed ISO-ENSO relationship.

4. Interannual variability and predictability

4.1 Causes of the year-to-year fluctuations

Webster (2006) and Yang and Lau (2006) provided comprehensive reviews on the interannual variability of the Asian monsoon. A great number of the monsoon literature has documented the year-to-year variability in various Asian monsoon regions, including the Indian monsoon (e.g., Mooley and Parthasarathy, 1984; Shukla and Mooley, 1987), the Indonesian monsoon (e.g., Yasunari and Suppiah, 1988; Hamada et al., 2002), the East Asian monsoon (e.g., Nitta, 1987; Huang and Wu, 1989; Li and Zeng, 2005; Zhou and Yu, 2005), and the WNP monsoon (e.g., Wu and Wang, 2000). The rainfall and circulation anomalies in many of the aforementioned regions exhibit a major 2-3-yr spectral peak (e.g., Meehl, 1987; Lau and Shen, 1988; Meehl and Ablaster, 2002). This is often referred to as the Tropospheric Biennial Oscillation (TBO) (Meehl, 1993).

Efforts have been made toward understanding the broad-scale interannual variability of the Asian-Australian monsoon (A-AM) system (e.g., Webster and Yang, 1992; Navarra et al., 1999; Miyakoda et al., 1999; Wang et al., 2003; Lau and Wang, 2006). Specific attention has been paid to the relationship between the ENSO and A-AM. Two principal modes of A-AM year-to-year variability have been identified (Wang et al., 2003, 2008c). The leading mode exhibits a prominent biennial tendency concurrent with the turnabout of ENSO, providing a new perspective of the seasonally evolving spatial-temporal structure for the TBO (Li et al., 2006). The second mode leads ENSO by 1 yr and is driven by SST anomalies associated with La Niña (Zhou et al., 2007).

The root causes of the interannual variation of the A-AM system have been extensively investigated over the past decade. The remote forcing from the ENSO through atmospheric teleconnection is no doubt an essential cause. But this is not the full story. For instance, the Indian monsoon rainfall variability appears to be determined by both the ENSO and equatorial Indian Ocean variability (Gadgil et al., 2004).

Since the SST anomalies in the monsoon ocean are, to a large extent, results of monsoon forcing, the monsoon-warm ocean interaction, rather than the warm ocean SST anomalies should be recognized as an essential process that determines the A-AM variability. The Indian-Ocean dipole (IOD) or zonal mode (Saji et al., 1999; Webster et al., 1999) is a beautiful example of the monsoon-ocean interaction. The IOD has impacts on monsoon rainfall anomalies in the Indian Ocean and over India and South Asia, East Africa, Maritime continent, East Asia, and the WNP (e.g., Saji et al., 1999; Guan and Yamagata, 2003). The processes supporting IOD have been attributed to the equatorial Bjerkness positive feedback (Webster et al., 1999; Saji et al., 1999). The WNP anomalous anticyclone that affects East Asian summer monsoon is another persuasive example (Wang et al., 2000; Lau and Wang, 2006). The convectively coupled Rossby wave-SST feedback can maintain both the WNP anticyclone and SST anomaly, which provides a prolonged impact of ENSO on the East Asian summer monsoon even when the SST anomalies in the eastern Pacific disappears. Monsoon-ocean interaction can also provide an important negative feedback through monsoon-induced anomalies in the surface heat fluxes (Lau and Nath, 2000) or through the Ekman transport of ocean heat (Webster et al., 2002; Loschnigg et al., 2003). This negative feedback often offsets the impacts of the remote ENSO forcing, making the Indian summer monsoon more resilient to interannual variation and more difficult to predict. In addition, this negative feedback is potentially important for supporting the monsoon TBO (Webster et al., 2002).

There is a hidden factor that contributes to the A-AM variation but has often been neglected. The monsoon basic flow not only regulates the nature of atmosphere-ocean interaction (Nicholls, 1983), but also significantly modifies the monsoon response to

remote ENSO forcing (Wang et al., 2003). Thus, the ENSO forcing, the monsoon-warm pool ocean interaction, and the influence of the annual cycle of the basic monsoon flow are three fundamental factors for understanding the behavior of the leading modes of A-AM variability (Wang et al., 2003).

The effects of the soil moisture and snow cover have long been recognized as sources of Asian monsoon variability, especially for the rainfall over the continental monsoon regions (see Yasunari (2006) for a detailed review). Yasunari (1991) and Dirmeyer et al. (1999) pointed out that the land surface conditions in spring have an impact on the following summer monsoon. The spring snow cover has shown to affect summer monsoon (e.g., Bamzai and Shukla, 1999; Zhang et al., 2004). But the role of the Eurasian snowfall does not overwhelm the SST impacts (Shen et al., 1998) and the snowfall itself may be affected by ENSO (Yang, 1996). The influence of snow cover and related soil moisture anomaly on the temperature and circulation anomalies in the lower troposphere may be limited to when and where the snow cover exists seasonally (Robock et al., 2003). More studies are needed to understand how snow in the Eurasian continent and/or over the Tibetan Plateau can affect succeeding monsoon activity.

In summary, our current understanding of the causes of the Asian monsoon interannual variability suggests that modeling and prediction of monsoon requires coupled atmosphere-ocean-land models; and an accurate forecast of the Asian monsoon requires realistic modeling of the ENSO, the annual cycle, the teleconnections associated with ENSO, the warm ocean-monsoon interaction, and the atmosphere-land surface interaction. These requirements make the modeling and dynamic prediction of Asian monsoon extremely challenge.

4.2 Modeling of monsoon interannual variations

Intercomparison of the capability of AGCMs to reproduce interannual monsoon anomalies has been one of the major efforts in identifying common problems of the models. Sperber and Palmer (1996) examined 32 AGCMs that participated in the AMIP project. They found that the Webster and Yang (1992) index (vertical wind shear) was better simulated than the all-Indian rainfall and that interannual variation was better simulated in the models which were able to generate a better climatology.

In assessment of the performance of 10 AGCMs that participated in the CLIVAR/AAMP AGCM intercomparison project, Wang et al. (2004) found that the models' poor simulation of anomalous summer monsoon rainfall is mainly due to lack of skill over Southeast Asia and the WNP region (5°-30°N, 80°-150°E). They proposed that the neglect of air-sea interaction is a major cause of the AMIP-type of models' failure. Examination of 5 AGCM-alone 21-yr ensemble simulations confirms that the AGCMs, when forced by observed SST, are unable to simulate Asian-Pacific summer monsoon rainfall (Wang et al., 2005b). The models tend to yield positive SST-rainfall correlations in the summer monsoon region that are at odds with observations. The observed negative SST-rainfall relationship (and rainfall leads SST variation) indicates atmosphere plays an active, not passive role. Thus, treating the atmosphere as a slave to specified SSTs may prohibit accurate simulation of summer monsoon rainfall anomalies.

Development of an effective strategy for improving models requires better design of the metrics for quantitative assessment of model performances and identification of the key modeling issues.

4.3 Predictability and seasonal prediction skills

In the past two decades, climate scientists have made ground-breaking progress in dynamic seasonal prediction. The advent of dynamic climate prediction can be traced back to El Niño forecast that used an intermediate-complexity coupled ocean-atmosphere model (Cane et al., 1986). Two types of prediction systems have been developed since the early 1990s. Bengtsson et al. (1993) proposed a "two-tier" approach for dynamical seasonal forecast, in which the global SST anomalies are first predicted, and an AGCM is subsequently forced by the pre-forecasted SST to make a future seasonal prediction. While the two-tier approach was a useful strategy to capture

better teleconnection, recent research advances suggest that prediction of certain phenomena (e.g., summer monsoon precipitation) requires taking into account local monsoon-ocean interactions (Wang et al., 2004; Wu and Kirtman, 2005; Kumar et al., 2005). Toward the end of the 20th century, a new era of seasonal forecast with CGCMs (also known as the one-tier approach) began, due to rapid progress made in coupled climate models (Latif et al., 2001; Davey et al., 2002) and due to a concerted international effort (through the Tropical Ocean-Global Atmosphere program) to monitor tropical ocean variations. Although the CGCMs still have significant systematic errors, many have demonstrated their capacity to reproduce realistic characteristics of ENSO (e.g., Latif et al., 1994; Ji et al., 1994). It has been increasingly recognized that the CGCMs are the ultimate tools for seasonal prediction.

For the two-tier systems, the physical basis for seasonal prediction lies in slowly varying lower boundary forcing, especially the anomalous SST (as well as the land surface) forcing (Charney and Shukla, 1981; Shukla, 1998). For the one-tier systems, prediction of ENSO and associated climate variability is essentially an initial value problem (Palmer et al., 2004). The slowly varying lower boundary of the atmosphere is evolving as a result of feedback from the atmosphere. The climate predictability in nature and in CGCMs comes from "slow" coupled (atmosphere-ocean-land-ice) dynamics and initial memories in ocean and land surfaces (Palmer et al., 2004; Wang et al., 2008b). Pertaining to the Asian monsoon, we need to find out how predictable the A-AM interannual variability is, especially in the continental monsoon region.

There are two types of error sources of dynamic seasonal prediction: atmospheric chaotic dynamics and uncertainties in the model parameterizations of unresolved sub-grid scale processes (Shukla et al., 2000; Kang and Shukla, 2006). Since the seasonal predictability does not depend on atmospheric initial conditions, an ensemble forecast technique with different atmospheric initial conditions was developed to reduce the errors arising from atmospheric chaotic dynamics. To alleviate the uncertainty arising from the sub-grid scales, stochastic physics schemes have been developed for individual models (Buizza et al., 1999). Meanwhile, a more effective way, the multi-model ensemble (MME) approach, was designed to reduce and quantify forecast uncertainties due to model formulation (Krishnamurti et al., 1999, 2000; Doblas-Reyes et al., 2000; Palmer, 2000). The idea behind the MME lies in that if the parameterization schemes in a group of models are independent of each other, the model errors due to the parameterization schemes may be random in nature; thus, an average approach may cancel out, at least partially, these errors.

It has firmly established that MME prediction is superior to the predictions made by any single-model component (Krishnamurti et al., 2000; Barnston et al., 2003; Doblas-Reyes et al., 2005). Based on this, a number of international projects have organized multi-model intercomparison and synthesis, among which the most comprehensive ones are the European Union-sponsored "Development of a European Multi-model Ensemble System for Seasonal to Inter-Annual Prediction (DEMETER; Palmer et al., 2004) and the Climate Prediction and its Application to Society (CliPAS) sponsored by the Asian-Pacific Economic Cooperation (APEC) Climate Center (APCC) (Wang et al., 2008a).

Recent evaluation of a 21-yr hindcast of 10 coupled climate models from the DEMETER and CliPAS shows that the one-month lead predictions capture the first two leading modes of variability with a verisimilitude that is at least comparable to or even better than that of the ERA-40 and NCEP-2 reanalyses (Wang et al., 2008a). This is due to the use of the multiple models and due to the fact that models include ocean-atmosphere interaction processes.

Wang et al. (2008b) assessed the current status of MME deterministic and probabilistic seasonal prediction based on 25-yr (1980–2004) retrospective forecasts performed by 14 APCC/CliPAS climate model systems (7 one-tier and 7 two-tier systems) and 7 DEMETER models' MME for the period of 1981-2001. Figure 4 shows that the forecast of monsoon precipitation remains a major challenge; and the seasonal rainfall predictions over land and during



Fig.4. Temporal correlation coefficients for precipitation between observation and one-month lead seasonal prediction for 1981-2003 obtained from 14 APCC/CliPAS models' MME system in (a) JJA and (b) DJF, respectively. The thin (thick) solid contours represent statistical significance of the correlation coefficients at 0.05 (0.01) confidence level. Adopted from Wang et al. (2008b).

local summer have little skill. The precipitation forecast skill averaged between 30°S and 30°N decreases away from the central-eastern Pacific, with the highest mean skill exceeding 0.5 found near the dateline (150°E–170°W) in JJA and from 150°E to 140°W in DJF; in contrast, the lowest skill is found over the tropical Africa.

5. Prospects on monsoon climate prediction

5.1 Seasonal prediction

Given the current levels of the climate models, a critical question is how we can get the best forecast through the MME. It has been speculated that the highest MME skill may be achievable by an optimal choice of a subgroup of models, drawing upon an individual model's skill and the mutual independence among the chosen models (Wang et al., 2008b).

While the MME is a proven approach that provides superior skill over any individual model, the skills of MMEs depend on having good models. Thus, improvement of individual models is of central importance for better climate forecasts. Then, what are the priorities we should take to improve climate models' physics?

The foremost factor leading to successful monsoon seasonal prediction is the model's capability to accurately forecast the amplitude, spatial pattern, and detailed evolution of ENSO cycle, because ENSO is the primary source for the monsoon and global climate predictability. This is particular true for a long lead seasonal forecast, because as forecast lead time increases, the model forecast tends to be determined by the model ENSO behavior (Jin et al., 2008). Therefore, Continuing improvement of the slow coupled dynamics in reproducing a realistic ENSO mode is a key for long-lead seasonal forecast.

The differences in the forecast skills over land areas between the CliPAS and DEMETER MMEs indicate potentials for further improvement of predictability over land (Wang et al., 2008b). The fact that MME has little skill in predicting precipitation over the continental region and during local summer season suggests potential importance of atmosphere-land interaction, because in these models the land surface initializations have considerable errors. There are urgent needs to (a) assess the impact of land surface initialization on the skill of seasonal and monthly forecast using a multi-model framework, (b) understand the roles of land surface processes and atmosphere-land interaction play in short-lead monthly and seasonal prediction, and (c) know how to improve seasonal prediction in continental monsoon regions.

Since the teleconnection both within the tropics and between the tropics and extratropics is a major source of predictability for the region outside of the eastern tropical Pacific, and since teleconnection is sensitive to mean climatology, continuing improvement of the mean state and seasonal cycle as well as statistical behavior of the transient atmospheric

circulation in coupled models is also of importance (Wang et al., 2008b). However, to what extent seasonal predictions depend on nonlinear rectification of high-frequency atmospheric and oceanic processes (so-called "noises") is not well known.

In the current CGCMs, lack of land surface initialization and use of fixed sea ice are common. Since the primary memory affecting slow coupled dynamics is stored in ocean subsurface layers (Rosati et al., 1997) and land surfaces, continuing improvement of coupled model initialization is an urgent task. In addition, accurate description of the warm pool ocean-monsoon interaction is another key issue.

Inclusion of anthropogenic (especially aerosols) and natural forcing (solar, volcanic, and aerosol) and a better representation of sea-ice may also benefit accurate seasonal forecast.

5.2 Intraseasonal prediction

Prediction of the occurrence of large-scale droughts and floods as ahead as possible is of considerable practical values for a wide range of interests, especially for the management of environmental and financial resources. The MJO and MISO strongly modulate occurrence of the tropical cyclones and floods and droughts. Thus, prediction of MJO and MISO has great potential to meet the societal demand.

First, future improvement of the intraseasonal forecast relies on improved climate models. Numerical experiments with the global cloud-system resolving model (GCRM) developed at the Frontier Research Center for Global Change (FRCGC) have demonstrated the promise of very high resolution modeling in simulating multi-scale structure of the MJO (Nasuno et al., 2007). Using the GCRM that allows direct coupling of the atmospheric circulation and clouds, Miura et al. (2007) successfully simulated the slow eastward migration of an MJO event during December 2006 to January 2007 (Fig.5). The results demonstrate the potential making of month-long MJO predictions when global cloud-resolving models with realistic initial conditions are used.



Fig.5. Time-longitude (Hovmöller) sections: precipitation (in m m h^{-1}) averaged between 10°S and 5°N from (a) the TRMM 3B42 data and (b) the 7-km grid run. The vorticity (10^{-5} s^{-1}) at the 850-hPa level averaged between 10° and 5°S from (c) the NCEP analyses and (d) the 7-km grid run. Adopted from Miura et al. (2007).

Very high-resolution models are necessary and critical for improved prediction of precipitation and statistical behavior of extreme events and high-impact weather. High resolution models may also be able to resolve the Meiyu front better. Synoptic disturbances play an important role in determination of extratropical monsoon climate variability. However, it remains to be demonstrated whether increased resolution and improved simulation of high-frequency perturbations would improve the slow coupled dynamics in the coupled climate models.

A key to improvement of intraseasonal prediction is to use coupled models and to better initialize the coupled system, because the intraseasonal prediction is sensitive to both atmospheric initial conditions (Waliser et al., 2003a; Reichler and Roads, 2005) and atmosphere-ocean interaction (Fu et al., 2007, 2008).

It should be addressed whether the MME approach could improve the skill of MJO/MISO prediction, although the MME has been proven to outperform a single model for the medium-range weather and hurricane forecasts (Krishnamurti et al., 1999, 2000). Application of MME to operational centers remains to be implemented. Building on experimental real-time efforts at CDC/NOAA (Waliser et al., 2006) and emerging operational efforts at CPC/NOAA and a number of other forecast centers, the US CLIVAR MJO Working Group has fostered the development of a multi-model operational MJO prediction framework. The most critical items needed to produce an MME forecast are: 1) an MJO-specific hindcast data set and 2) the development of an MME methodology relevant to the MJO.

Looking forward, the increase in the new observations and the advances in cloud resolving models, computer power, and communication technology are expected to provide breakthroughs in monsoon climate prediction.

Acknowledgements. I acknowledge numerous stimulating discussions that have fostered the present review from my collaborators and colleagues at the IPRC, the APCC Science Advisory Committee, the AMY Science Steering Committee, the CLIVAR AAMP, the WCRP/IMS Scoping Group, and the US CLIVAR/MJO working group; in particular Drs. X. Fu, J.-Y. Lee, Q. Ding, K. Kikuchi, D. E. Waliser, J. Matsumoto, J. Shukla, In-Sik Kang, H. H. Hendon, C. P. Chang, T. Yasunari, G.-X. Wu, B. Kirtman, A. Kumar, and S. Schubert. Thanks also go to Y. Ding and Z. Liu for their kind invitation and Z. Wu for assistance in compilation of the reference list.

REFERENCES

Annamalai, H., and J. Slingo, 2001: Active/break cycles: Diagnosis of the intraseasonal variability of the Asian summer monsoon. *Climate Dyn.*, **18**, 85–102.

Bamzai, A., and J. Shukla, 1999: Relation between Eurasian snow cover, snow depth and the Indian summer monsoon: An observational study. *J. Climate*, **12**, 3117–3132.

Barnston, A. G., S. J. Mason, L. Goddard, D. G. Dewitt, and S. E. Zebiak, 2003: Multimodel ensembling in seasonal climate forecasting at IRI. *Bull. Amer. Meteor. Soc.*, **84**, 1783–1796.

Bengtsson, L., U. Schlese, E. Roeckner, M. Latif, T. P. Barnett, and N. E. Graham, 1993: A two-tiered approach to long-range climate forecasting. *Science*, **261**, 1027–1029.

Buizza, R., M. Miller, and T. N. Palmer, 1999: Stochastic representation of model uncertainties in the ECMWF ensemble prediction system. *Quart. J. Roy. Meteor. Soc.*, **125**, 2887–2908.

Cane, M. A., S. E. Zebiak, and S. C. Dolan, 1986: Experimental forecasts of El Niño. *Nature*, **321**, 827–832.

Chang, C.-P., B. Wang, and N.-C. G. Lau, 2005: The global monsoon system: research and forecast. WMO/TD No. 126, Secretariat of WMO, Geneva, Switzerland.

Chang, C.-P., Z. Wang, J. McBride, and C. H. Liu, 2005: Annual cycle of Southeast Asia-Maritime Continent rainfall and the asymmetric monsoon transition. *J. Climate*, **18**, 287–301.

Chang, C.-P., Z. Wang, and H. Hendon, 2006: The Asian winter monsoon. *The Asian Monsoon*. Wang B., Ed., Praxis, Berlin, 89–127.

Charney, J., and J. Shukla, 1981: Predictability of monsoons. *Monsoon Dynamics*. Lighthill J., and R. P. Pearce, Eds., Cambridge University Press, Cambridge, UK, 99–108.

Chen, T., and J. Chen, 1993: The 10-20-day mode of the

1979 Indian monsoon: its relation with the time variation of monsoon rainfall. *Mon. Wea. Rev.*, **121**, 2465–2482.

Davey, M. K., and Coauthors, 2002: STOIC: A study of coupled GCM climatology and variability in tropical ocean regions. *Climate Dyn.*, **18**, 403–420.

Ding, Q., and B. Wang, 2007: Intraseasonal teleconnection between the Eurasian wave train and Indian summer monsoon. *J. Climate*, **20**, 3751–3767.

Ding, Y., 1992: Summer monsoon rainfalls in China. *J. Meteor. Soc. Japan*, **70**, 397–421.

Ding, Y., 1994: *Monsoons Over China.* Springer, New York, 419.

Ding, Y., 2004: Seasonal march of the East-Asian summer monsoon. *The East Asian Monsoon.* C.-P. Chang Ed., World Scientific Publishing, 3–53.

Dirmeyer, P., A. Dolman, and N. Sato, 1999: The pilot phase of the global soil wetness project. *Bull. Amer. Meteor. Soc.*, **80**, 851–878.

Doblas-Reyes, F. J., M. Déqué, and J.-P. Piedelievre, 2000: Model and multimodel spread and probabilistic seasonal forecasts in PROVOST. *Quart. J. Roy. Meteor. Soc.*, **126**, 2069–2088.

Doblas-Reyes, F. J., R. Hagedorn, and T. N. Palmer, 2005: The rationale behind the success of multimodel ensembles in seasonal forecasting. Part II: Calibration and combination. *Tellus A,* **57**, 234–252.

Drbohlav, H., and B. Wang, 2005: Mechanism of the northward propagating intraseasonal oscillation in the South Asian monsoon region: Results from a zonally averaged model. *J. Climate*, **18**, 952–972.

Fu, X., B. Wang, T. Li, and J. P. McCreary, 2003: Coupling between northward propagating, intraseasonal oscillations and sea-surface temperature in the Indian Ocean. *J. Atmos. Sci.,* **60**, 1733–1753.

Fu, X., B. Wang, D. E. Waliser, and L. Tao, 2007: Impact of atmosphere-ocean coupling on the predictability of monsoon intraseasonal oscillations. *J. Atmos. Sci.*, **64**, 157–174.

Fu, X., B. Wang, Q. Bao, P. Liu, and B. Yang, 2008: Experimental dynamical forecast of an MJO event observed during TOGA-COARE period. *Atmos. Oceanic Sci. Lett.*, in press.

Gadgil, S., P. Vinayachandran, P. Francis, and S. Gadgil, 2004: Extremes of the Indian summer monsoon rainfall, ENSO and equatorial Indian Ocean oscillation. *Geophys. Res. Lett.*, **31**, L12213, doi: 10.1029/2004GL019733.

Goswami, B., and J. Shukla, 1984: Quasi-periodic oscillations in a symmetric general circulation model. *J. Atmos. Sci.,* **41**, 20–37.

Guan, Z., and T. Yamagata, 2003: The unusual summer of 1994 in East Asia: IOD teleconnections. *Geophys. Res. Lett.,* **30**(10), 1544, doi: 10.1029/2002GL016831.

Hamada, J.-I., M. Yamanaka, J. Matsumoto, S. Fukao, P. Winarso, and T. Sribimawati, 2002: Spatial and temporal variations of the rainy season over Indonesia and their link to ENSO. *J. Meteor. Soc. Japan*, **80**, 285–310.

Hendon, H. H., B. Liebmann, M. Newman, J. D. Glick, and J. Schemm, 1999: Medium range forecast errors associated with active episodes of the MJO. *Mon. Wea. Rev.*, **128**, 69–86.

Hendon, H., M. Wheeler, and C. Zhang, 2007: Seasonal dependence of the MJO-ENSO relationship. *J. Climate*, **20**, 531–543.

Hsu, H.-H., and C.-H. Weng, 2001: Northwestward propagation of the intraseasonal oscillation during the boreal summer: mechanism and structure. *J. Climate*, **14**, 3834–3850.

Hsu, H.-H., C.-H. Weng, and C.-H. Wu, 2004: Contrasting characteristics between the northward and eastward propagation of the intraseasonal oscillation during the boreal summer. *J. Climate*, **17**, 727–743.

Hsu, H.-H., and M.-Y. Lee, 2005: Topographic effects on the eastward propagation and initiation of the Madden-Julian Oscillation. *J. Climate*, **18**, 795–809.

Huang, R., and Y. Wu, 1989: The influence of ENSO on the summer climate change in China and its mechanism. *Adv. Atmos. Sci.*, **6**, 21–32.

Hung, C.-W., X. Liu, and M. Yanai, 2004: Symmetry and asymmetry of the Asian and Australian summer monsoon. *J. Climate*, **17,** 2413–2426.

Ji, M., A. Kumar, and A. Leetmaa, 1994: An experimental coupled forecast system at the National Meteorological Center: Some early results. *Tellus*, **46**A, 398–418.

Jiang, X., and T. Li, 2005: Reinitiation of the boreal summer intraseasonal oscillation in the tropical Indian Ocean. *J. Climate*, **18**, 3777–3795.

Jiang, X., T. Li, and B. Wang, 2004: Structures and mechanisms of the northward propagating boreal summer intraseasonal oscillation. *J. Climate*, **17**, 1022–1039.

Jin, E. K, J. L. Kinter III, B. Wang, and Co authors, 2008: Current status of ENSO prediction skill in coupled ocean-atmosphere model. *Climate Dyn.*, doi:10.1007/s00382-008-0418-2.

Jones, C., D. E. Waliser, J.-K. E. Schemm, and W. K.-M. Lau, 2000: Prediction skill of the Madden-Julian Oscillation in dynamical extended range forecasts. *Climate Dyn.*, **16**, 273–289.

Kang, I.-S., and J. Shukla, 2006: Dynamic seasonal prediction and predictability (Chap.15). *The Asian Monsoon*. Springer Praxis, Chicherster, 585–612.

Kang, I.-S., K. Jin, B. Wang, K.-M. Lau, J. Shukla, and Coauthors, 2002: Intercomparison of the climatological variations of Asian summer monsoon precipitation simulated by 10 GCMs. *Climate Dyn.*, **19**, 383–395.

Kawamura, R., T. Murakami, and B. Wang, 1996: Tropical and midlatitude 45-day perturbations during the northern summer. *J. Meteor. Soc. Japan*, **74**, 867–890.

Kemball-Cook, S., and B. Wang, 2001: Equatorial waves and air-sea interaction in the boreal summer intraseasonal oscillation. *J. Climate*, **14**, 2923–2942.

Kiguchi, M., and J. Matsumoto, 2005: The rainfall phenomena during the pre-monsoon period over the Indochina peninsula in the GAME-IOP year. *J. Meteor. Soc. Japan*, **83**, 89–106.

Kikuchi, K., and B. Wang, 2008: Diurnal precipitation regimes in the tropics. *J. Climate*, in press.

Kim, H.-M, I.-S. Kang, B. Wang, and J.-Y. Lee, 2008: Interannual variations of the Boreal summer intraseasonal variability predicted by ten atmosphere-ocean coupled models. *Climate Dyn.*, **30**, 485-496, doi: 10.1007/s00382-007-0292-3.

Krishnamurthy, V., and J. Shukla, 2008: Seasonal persistence and propagation of intraseasonal patterns over the Indian monsoon region. *Climate Dyn.*, **30**, 353–369.

Krishnamurti, T., and H. Bhalme, 1976: Oscillations of a monsoon system. Part I. Observational aspects. *J. Atmos. Sci.,* **33**, 1937–1954.

Krishnamurti, T. N., C. M. Kishtawal, T. E. LaRow, D. R. Bachiochi, Z. Zhang, C. E. Williford, S. Gadgil, and S. Surendran, 1999: Improved skills for weather and seasonal climate forecasts from multi-model superensemble. *Science*, **285**, 1548–1550.

Krishnamurti, T. N., C. M. Kishtawal, Z. Zhang, T. E. LaRow, D. R. Bachiochi, Z. Zhang, C. E. Williford, S. Gadgil, and S. Sajani, 2000: Multimodel ensemble forecasts for weather and seasonal climate. *J. Climate*, **13**, 4196–4216.

Kumar, K. K., M. Hoerling, and B. Rajagopalan, 2005: Advancing dynamical prediction of Indian monsoon rainfall, *Geophys. Res. Lett.*, **32**, L08704, doi:10.1029/2004GL021979.

Latif, M., T. P. Barnett, M. A. Cane, M. Flugel, N. E. Graham, H. von Storch, et al., 1994: A review of ENSO prediction studies. *Clim. Dyn.*, **9**, 167–179, doi:10.1007/BF00208250.

Latif, M., and Coauthors, 2001: ENSIP: The El Niño simulation intercomparison project. *Climate Dyn.*, **18**, 255–276.

Lau, K.-M., and P. Chan, 1986: Aspects of the 40-50 day oscillation during northern summer as inferred from OLR. *Mon. Wea. Rev.,* **114**, 1354–1367.

Lau, K.-M., M. Kim, and K. Kim, 2006: Aerosol induced anomalies in the Asian summer monsoon−the role of the Tibetan Plateau. *Climate Dyn.*, **26**, 855–864.

Lau, K. M., J.-Y. Lee, K.-M. Kim, and I.-S. Kang, 2004: The North Pacific as a regulator of summertime climate over Eurasia and North America. *J. Climate*, **17**, 819–833.

Lau, K.-M., and S. Shen, 1988: On the dynamics of intraseasonal oscillations and ENSO. *J. Atmos. Sci.*, **45**, 1781–1797.

Lau, K.-M., and S. Yang, 1997: Climatology and interannual variability of the Southeast Asian summer monsoon. *Adv. Atmos. Sci.*, **14**, 141–162.

Lau, N.-C., and M. J. Nath, 2000: Impact of ENSO on the variability of the Asian-Australian monsoons as simulated in GCM experiments. *J. Climate*, **13**, 4287–4309.

Lau, N.-C., and B. Wang, 2006: Interactions between the Asian monsoon and the El Niño/Southern Oscillation. *The Asian Monsoon.* Wang B., Ed. Springer/Praxis Publishing Ltd., New York, 479–511.

Lawrence, D., and P. Webster, 2002: The boreal summer intraseasonal oscillation: Relationship between northward and eastward movement of convection. *J. Atmos. Sci.*, **59**, 1593–1606.

Lee, J.-Y., B. Wang, I.-S. Kang, J.-S. Kug, J. Shukla, E.-K. Jin, and Coauthors, 2008: Performance of climate prediction models on annual modes of precipitation and its relation with seasonal prediction. *Climate Dyn.*, in press.

Li, J., and Q. Zeng, 2002: A unified monsoon index. *Geophys. Res. Lett.*, **29**(8), 1274, doi:10.1029/2001GL013874.

Li, J., and Q. Zeng, 2005: A new monsoon index, its interannual variability and relation with monsoon precipitation. *Climate Environ. Res.*, **10**, 351–365.

Li, T., P. Liu, X. Fu, and B. Wang, 2006: Spatiotemporal structures and mechanisms of the tropospheric biennial oscillation in the Indo-Pacific warm ocean regions. *J. Climate*, **19**, 3070–3087.

Liess, S., D. E. Waliser, and S. D. Schubert, 2005: Predictability studies of the intraseasonal oscillation with the ECHAM5 GCM. *J. Atmos. Sci.*, **62**, 3320–3336.

LinHo, and B. Wang, 2002: The time-space structure of the Asian-Pacific summer monsoon: A fast annual cycle view. *J. Climate*, **15**, 2001–2019.

Lorenz, E. N., 1963: Deterministic nonperiodic flow. *J. Atmos. Sci.*, **20**, 130–141.

Loschnigg, J., G. Meehl, J. Webster, J. Arblaster, and G. Compo, 2003: The Asian monsoon, the tropospheric biennial oscillation, and the Indian Ocean zonal mode in the NCAR GCM. *J. Climate*, **15**, 2001–2019.

Madden, R. A., and P. R. Julian, 1971: Detection of a 40-50 day oscillation in the tropics. *J. Atmos. Sci.*, **43**, 3138–3158.

Madden, R. A., and P. R. Julian, 1972: Description of global-scale circulation cells in the tropics with a 40-50 day period. *J. Atmos. Sci.*, **29**, 1109–1123.

Maloney, E. D., and D. Hartmann, 1998: Frictional moisture convergence in a composite life cycle of the Madden-Julian Oscillation. *J. Climate*, **11**, 2387–2403.

Maloney, E. D., and D. L. Hartmann, 2001: The sensitivity of intraseasonal variability in the NCAR CCM3 to changes in convective parameterization. *J. Climate*, **14**, 2015–2034.

Matsumoto, J., 1992: The seasonal changes in Asian and Australian monsoon regions. *J. Meteor. Soc. Japan*, **70**, 257–273.

Matsumoto, J., 1997: Seasonal transition of summer rainy season over Indochina and adjacent monsoon region. *Adv. Atmos. Sci.,* **14**, 231–245.

Matsumoto, J., and T. Murakami, 2002: Seasonal migration of monsoons between the Northern and Southern Hemisphere as revealed from equatorially symmetric and asymmetric OLR data. *J. Meteor. Soc. Japan*, **80**, 419–437.

Meehl, G., 1987: The annual cycle and interannual variability in the tropical Pacific and Indian Ocean region. *Mon. Wea. Rev.*, **115**, 27–50.

Meehl, G., 1993: A coupled air-sea biennial mechanism in the tropical Indian and Pacific regions: Role of the ocean. *J. Climate*, **6**, 31–41.

Meehl, G., and J. Arblaster, 2002: The tropospheric biennial oscillation and Asian-Australian monsoon rainfall. *J. Climate*, **15**, 722–744.

Minoura, D., R. Kawamura, and T. Matsuura, 2003: A mechanism of the onset of the South Asian summer monsoon. *J. Meteor. Soc. Japan*, **81**, 563–580.

Miura, H., M. Satoh, T. Nasuno, A. T. Noda, and K. Oouchi, 2007: A Madden-Julian Oscillation event realistically simulated by a global cloud-resolving model. *Science*, **318**, doi: 10.1126/science.1148443.

Miyakoda, K., A. Navarra, and M. N. Ward, 1999: Tropical-wide teleconnection and oscillation. II: The ENSO-monsoon system. *Quart. J. Roy. Meteor. Soc.*, **125**, 2937–2963.

Mooley, D., and B. Parthasarathy, 1984: Fluctuation in all-India summer monsoon rainfall during 1871–1985. *Climate Change,* **6**, 287–301.

Mori, S., J.-I. Hamada, Y. Tauhid, M. Yamanaka, N. Okamoto, F. Murata, N. Sakurai, H. Hashiguchi, and T. Sribimawati, 2004: Diurnal land-sea rainfall peak migration over Sumatera Island, Indonesian maritime continent, observed by TRMM satellite and intensive rawinsonde soundings. *Mon. Wea. Rev.*, **132**, 2021–2039.

Murakami, M., 1983: Analysis of the deep convective activity over the western Pacific and Southeast Asia. Part I: Diunal variation. *J. Meteor. Soc. Japan*, **61**, 60–76.

Nakazawa, T., 1992: Seasonal phase lock of intraseasonal variation during the Asian summer monsoon. *J. Meteor. Soc. Japan,* **70**, 257–273.

Nasuno, T., H. Tomita, S. Iga, H. Miura, and M. Satoh, 2007: Multiscale organization of convection simulated with explicit cloud processes on an aquaplanet. *J. Atmos. Sci.*, **64**, 1902–1921.

Navarra, A., M. Ward, and K. Miyakoda, 1999: Tropical-wide teleconnection and oscillation. I: Teleconnection indices and type I/type II states. *Quart. J. Roy. Meteor. Soc.*, **125**, 2909–2935.

Nesbitt, S. W., and E. J. Zipser, 2003: The diurnal cycle of rainfall and convective intensity according to three years of TRMM measurements. *J. Climate*, **16**(10), 1456-1475.

Nicholls, N., 1983: Air-sea interaction and the quasi-biennial oscillation. *Mon. Wea. Rev.*, **106**, 1505–1508.

Nitta, T., 1987: Convective activities in the tropical western Pacific and their impact on the Northern Hemisphere summer monsoon. *J. Meteor. Soc. Japan*, **65**, 373–390.

Nitta, T., and S. Sekine, 1994: Diurnal variation of convective activity over the tropical western Pacific. *J. Meteor. Soc. Japan*, **72**, 627–641.

Ohsawa, T., H. Ueda, T. Hayashi, A. Watanabe, and J. Matsumoto, 2001: Diurnal variation of convective activity and rainfall in tropical Asia. *J. Meteor. Soc. Japan*, **79**, 333–352.

Palmer, T. N., 1994: Chaos and predictability in forecasting the monsoons. *Proc. Indian Natl. Sci. Acad.*, **60**, 57–66.

Palmer, T. N., 2000: Predicting uncertainty in forecasts of weather and climate. *Rep. Prog. Phys.*, **63**, 71–116.

Palmer, T. N., A. Alessandri, U. Andersen, P. Cantelaube, M. Davey, and Coauthors, 2004: Development of a European multi-model ensemble system for seasonal to interannual prediction (DEMETER). *Bull. Amer. Meteor. Soc.*, **85**, 853–872.

Pegion, K., and B. P. Kirtman, 2008: The impact of air-sea interactions on the predictability of the tropical intraseasonal oscillation. *J. Climate*, in press.

Ramage, R., 1971: *Monsoon Meteorology.* Academic Press, New York, 296 pp.

Rao, Y. P., 1976: Southwest monsoon: Synoptic meteorology. Meteor. Monogr., 1, Indian Meteorological Department, 367 pp.

Reichler, T., and J. O. Roads, 2005: Long-range predictability in the tropics. Part II: 30-60-day variability. *J. Climate*, **18**, 634–650.

Robock, A., M. Mu, K. Vinnikov, and D. Robinson, 2003: Land surface conditions over Eurasia and Indian summer monsoon rainfall. *J. Geophy. Res.*, **108**(D4), 4131, doi:10.1029/2002JD002286.

Rosati, A., K. Miyakoda, and R. Gudgel, 1997: The impact of ocean initial conditions on ENSO forecasting with a coupled model. *Mon. Wea. Rev.*, **125**(5), 754–772.

Saji, N., B. Goswami, P. Vinayachandran, and T. Yamagata, 1999: A dipole mode in the tropical Indian Ocean. *Nature*, **401**, 360–363.

Seo, K. -H., J. -K. E. Schemm, C. Jones, and S. Moorthi, 2005: Forecast skill of the tropical intraseasonal oscillation in the NCEP GFS dynamical extended range forecasts. *Climate Dyn.*, **25**, 265–284.

Shen, X., M. Kimoto, and A. Sumi, 1998: Role of land surface processes associated with interannual variability of broad-scale Asian summer monsoon as simulated by the CCSR/NIES AGCM. *J. Meteor. Soc. Japan.*, **76**, 217–236.

Shukla, J., 1998: Predictability in the midst of chaos: A scientific basis for climate forecasting. *Science*, **282**, 728–731.

Shukla, J., J. Anderson, D. Baumhefner, C. Brankovic, Y. Chang, E. Kalnay, L. Marx, T. Palmer, D. Paolino, J. Ploshay, S. Schubert, D. Straus, M. Suarez, and J. Tribbia, 2000: Dynamical seasonal prediction. *Bull. Amer. Meteor. Soc.*, **81**, 2593–2606.

Shukla, J., and D. Mooley, 1987: Empirical prediction of the summer monsoon rainfall over India. *Mon. Wea. Rev.*, **115**, 695–704.

Sikka, D., and S. Gadgil, 1980: On the maximum cloud zone and the ITCZ over Indian longitudes during the southwest monsoon. *Mon. Wea. Rev.*, **108**, 1840–1853.

Slingo, J. M., K. R. Sperber, J. S. Boyle, et al., 1996: Intraseasonal oscillation in 15 atmospheric general circulation models: Results from an AMIP diagnostic subproject. *Climate Dyn.*, **12**, 325–357.

Sorooshian, S., X. Gao, K. Hsu, R. Madox, Y. Hong, and Coauthors, 2002: Diurnal variability of tropical rainfall retrieved from combined GOES and TRMM satellite information. *J. Climate*, **15**, 983–1001.

Sperber, K., and T. Palmer, 1996: Interannual tropical rainfall variability in general circulation model simulations associated with the atmospheric model intercomparison project. *J. Climate*, **9**, 2727–2750.

Sperber, K. R., J. Slingo, and H. Annamalai, 2000: Predictability and the relationship between subseasonal and interannual variability during the Asian summer monsoon. *Quart. J. Roy. Meteor. Soc.*, **126**, 2545–2574.

Sperber, K., and Coauthors, 2001: Dynamical seasonal predictability of the Asian summer monsoon. *Mon. Wea. Rev.*, **129**, 2226–2248.

Sperber, K. R., D. E. Waliser, and Coauthors, 2008: New approaches to understanding, simulating, and forecasting the Madden-Julian oscillation. *Bull. Amer. Meteor. Soc.*, **89**, submitted.

Sperber, K., and T. Yasunari, 2006: Workshop on monsoon climate systems. *Bull. Amer. Meteor. Soc.*, **87**, 1399–1403.

Stephens, G., P. Webster, R. Johnson, R. Engelen, and T. L'Ecuyer, 2004: Observational evidence for the mutual regulation of the tropical hydrological cycle and tropical sea surface temperatures. *J. Climate*, **17**, 2213–2224.

Tanaka, M., 1992: Intraseasonal oscillation and the onset and retreat dates of the summer monsoon over the east, southeast and western north Pacific region using GMS high cloud amount data. *J. Meteor. Soc. Japan*, **70**, 613–629.

Taniguchi, K., and T. Koike, 2007: Increasing atmospheric temperature in the upper troposphere and cumulus convection over the eastern part of the Tibetan Plateau in the pre-monsoon season of 2004. *J. Meteor. Soc. Japan*, **85**A, 271–294.

Tao, S., and L. Chen, 1987: A review of recent research on the East Asian summer monsoon in China. *Monsoon Meteorology*. C. -P. Chang and T. N. Krishnamurti, Eds., Oxford University Press, New York, 60–92.

Trenberth, K. E., D. P. Stepaniak, and J. M. Caron, 2000: The Global monsoon as seen through the divergent atmospheric circulation. *J. Climate*, **13**, 3969–3993.

Ueda, H., 2005: Air-sea coupled process involved in stepwise seasonal evolution of the Asian summer monsoon. *Geogr. Rev. Japan*, **86**, 825–841.

Ueda, H., and H. T. Yasunari, 1996: Maturing process of summer monsoon over the western North Pacific-A coupled ocean/atmosphere system. *J. Meteor. Soc. Japan*, **74**, 493–508.

Ueda, H., H. Kamahori, and N. Yamazaki, 2003: Seasonal contrasting features of heat and moisture budgets between the eastern and western Tibetan Plateau during the GAME IOP. *J. Climate*, **16**, 2309–2324.

Vitart, F., S. Woolnough, M. A. Balmaseda, and Coauthors, 2008: Monthly forecast of the Madden-Julian Oscillation using a coupled GCM. *Mon. Wea. Rev.*, **135**, 2700–2715.

Waliser, D. E., 2005: Predictability and forecasting tropical intraseasonal variability. *Intraseasonal Variability of the Atmosphere-Ocean Climate System*. Lau W. K. M., and D. E. Waliser, Eds., Springer, Heidelberg, Germany, 401–434.

Waliser, D. E., 2006: Intraseasonal variability. *The Asian Monsoon*. Wang, B. Ed., Springer/Praxis Publishing Ltd., New York, 203–259.

Waliser, D. E., K.-M. Lau, W. Stern, and C. Jones, 2003a: Potential predictability of the Madden-Julian Oscillation. *Bull. Amer. Meteor. Soc.*, **84**, 33–50.

Waliser, D. E., K. Jin, I.-S. Kang, W. Stern, S. Schubert, and Coauthors, 2003b: AGCM simulations of intraseasonal variability associated with the Asian summer monsoon. *Climate Dyn.*, **21**, 423–446.

Waliser, D. E., K. Weickmann, R. Dole, S. Schubert, O. Alves, and Coauthors, 2006: The experimental MJO prediction project. *Bull. Amer. Meteor. Soc.*, **87**, 425–431.

Wang, B., 1994: Climatic regimes of tropical convection and rainfall. *J. Climate*, **7**, 1109–1118.

Wang, B., 2005: Chapter 10 Theory. *Intraseasonal Variability of the Atmosphere-Ocean Climate System*. Lau W. K. M., and D. E. Waliser, Eds., Springer, Heidelberg, Germany, 401–434.

Wang, B., 2006: *The Asian Monsoon*. Springer-Verlag/Praxis Publishing, New York, 787 pp.

Wang, B., and Q. Ding, 2008: The global monsoon: Major modes of annual variations in the tropics. *Dynamics of Atmos. and Ocean*, special issue **2**, 165–183, doi: 10.1016/j.dynatmoce.2007.05.002.

Wang, B., and LinHo, 2002: Rainy seasons of the Asian-Pacific monsoon. *J. Climate*, **15**, 386–398.

Wang, B., and H. Rui, 1990: Dynamics of the coupled moist Kelvin-Rossby wave on an equatorial beta-plane. *J. Atmos. Sci.*, **47**, 397–413.

Wang, B., and X. Xie, 1997: A model for the boreal summer intraseasonal oscillation. *J. Atmos. Sci.*, **54**, 72–86.

Wang, B., and X. Xu, 1997: Northern Hemisphere summer monsoon singularities and climatological intraseasonal oscillation. *J. Climate*, **10**, 1071–1085.

Wang, B., I.-S. Kang, and J.-Y. Lee, 2004: Ensemble simulations of Asian-Australian monsoon variability by 11 AGCMs. *J. Climate*, **17**, 803–818.

Wang, B., P. J. Webster, and H. Teng, 2005a: Antecedents and self-induction of the active-break Indian summer monsoon. *Goephys. Res. Lett.*, **32**, L04704.

Wang, B., R. Wu, and X. Fu, 2000: Pacific-East Asian teleconnection: How does ENSO affect East Asian climate? *J. Climate*, **13**, 1517–1536.

Wang, B., R. Wu, and T. Li, 2003: Atmosphere-warm ocean interaction and its impact on Asian-Australian monsoon variation. *J. Climate*, **16**, 1195–1211.

Wang, B., Q. Ding, X. Fu, I.-S. Kang, K. Jin, J. Shukla, and F. Doblas-Reyes, 2005b: Fundamental challenges in simulation and prediction of summer monsoon rainfall. *Geophys. Res. Lett.*, **32**, L15711, doi: 10.1029/2005GL022734 12.

Wang, B., P. Webster, K. Kikuchi, T. Yasunari, and Y. Qi, 2006: Boreal summer quasi-monthly oscillation in the global tropics. *Climate Dyn.*, **27**, 661–675.

Wang, B., J.-Y. Lee, I.-S. Kang, J. Shukla, J.-S. Kug, A. Kumar, J. Schemm, J.-J. Luo, T. Yamagata, and C.-K. Park, 2008a: How accurately do coupled climate models predict the leading modes of Asian-Australian monsoon interannual variability? *Climate Dyn.*, **30**, 605–619. doi: 10.1007/s00382-007-0310-5.

Wang, B., J.-Y. Lee, J. Shukla, I.-S. Kang, C.-K. Park and Coauthors, 2008b: Advance and prospectus of seasonal prediction: Assessment of APCC/CliPAS 14-model ensemble retrospective seasonal prediction (1980–2004). *Climate Dyn.*, doi:10.1007/s00382-008-0460-0.

Wang, B., J. Yang, T. Zhou, and B. Wang, 2008c: Interdecadal changes in the major modes of Asian-Australian monsoon variability: strengthening relationship with ENSO since the late 1970s. *J. Climate*, **21**, 1771–1789, doi: 10.1175/2007JCLI 1981.1.

Webster, P. J., 1983: Mechanisms of monsoon low-frequency variability: Surface hydrological effects. *J. Atmos. Sci.*, **40**, 2110–2124.

Webster, P. J., 1987: The elementary monsoon. *Monsoons.* Fein J. S., and P. L. Stephens, Eds., John Wiley, New York, N.Y. 3–32.

Webster, P. J., 2006: The coupled monsoon system. *The Asian monsoon.* Wang B., Ed., Springer/Praxis Publishing Ltd., New York, 3–65.

Webster, P. J., and S. Yang, 1992: Monsoon and ENSO: Selectively interactive systems. *Quart. J. Roy. Meteor. Soc.*, **118**, 877–926.

Webster, P. J., E. F. Bradley, C. W. Fairall, J. S. Godfrey, P. Hacker, and Coauthors, 2002: The JASMINE pilot study. *Bull. Amer. Meteor. Soc.*, **83**, 1603–1630.

Webster, P. J., V. Magana, T. Palmer, J. Shukla, R. Tomas, M. Yanai, and T. Yasunari, 1998: Monsoons: Processes, predictability, and the prospects for prediction. *J. Geophys. Res.,* **103**, 14451–14510.

Webster, P. J., A. Moore, J. Loschnigg, and R. Leben, 1999: Coupled ocean-atmosphere dynamics in the Indian Ocean during 1997-1998. *Nature*, **401**, 356–360.

Wen, M., and R. Zhang, 2008: Quasi-biweekly oscillation of the convection around Sumatra and low-level tropical circulation in boreal spring. *Mon. Wea. Rev.*, **136**, 189–205.

Wu, G., and Y. Zhang, 1998: Tibetan Plateau forcing and the timing of the monsoon onset over South Asia and the South China Sea. *Mon. Wea. Rev.*, **126**, 913–927.

Wu, R., and B. Wang, 2000: Interannual variability of summer monsoon onset over the western North Pacific and the underlying processes. *J. Climate*, **13**, 2483–2501.

Wu, R., and B. Wang, 2001: Multi-stage onset of summer monsoon over the western North Pacific. *Climate Dyn.*, **17**, 277–289.

Wu, R., and B. Kirtman, 2005: Roles of Indian and Pacific Ocean air-sea coupling in tropical atmospheric variability. *Clim. Dyn.* **25**, 155–170.

Wu, Z., and J. Li, 2008: Prediction of the Asian-Australian monsoon interannual variations with the grid-point atmospheric model of IAP LASG (GAMIL). *Adv. Atmos. Sci.*, **25**(3), 387–394.

Yang, S., 1996: ENSO-snow-monsoon association and seasonal-interannual prediction. *Int. J. Climatol.*, **16**, 125–134.

Yang, S., and K.-M. Lau, 2006: Interannual variability of the Asian monsoon. *The Asian monsoon.* Wang B., Ed., Springer/Praxis Publishing Ltd., New York, 259–293.

Yang, G., and J. Slingo, 2001: The diurnal cycle in the tropics. *Mon. Wea. Rev.*, **129**, 784–801.

Yasunari, T., 1979: Cloudiness fluctuations associated with the Northern Hemisphere summer monsoon. *J. Meteor. Soc. Japan*, **57**, 227–242.

Yasunari, T., 1980: A quasi-stationary appearance of 30-40 day period in the cloudiness fluctuations during the summer monsoon over India. *J. Meteor. Soc. Japan*, **58**, 225–229.

Yasunari, T., 1991: The monsoon year: A new concept of the climate year in the tropics. *Bull. Amer. Meteor. Soc.*, **72**, 1331–1338.

Yasunari, T., 2006: Atmosphere-land interaction. *The Asian Monsoon.* Wang B., Ed., Springer/Praxis Publishing, New York, 459–478.

Yasunari, T., and R. Suppiah, 1988: Some problems on the interannual variability of Indonesian monsoon rainfall. *Tropical Rainfall Measurements.* Theon J. S., and N. Fugono, Eds., Deepak, Hampton, 113–122.

Yu, R., T. Zhou, A. Xiong, Y. Zhu, and J. Li, 2007: Diurnal variations of summer precipitation over contiguous China. *Geophys. Res. Lett.*, **34**, L01704, doi:10.1029/2006GL028129.

Zeng, Q., and J. Li, 2002: On the interaction between Northern and Southern Hemispheric atmospheres and the essence of tropical monsoon. *Chin. J. Atmos. Sci.*, **26**, 207–226.

Zhang, Y., T. Li, and B. Wang, 2004: Decadal change of the spring snow depth over Tibetan Plateau: The associated circulation and influence on the East Asian summer monsoon. *J. Climate*, **17**, 2780–2993.

Zhou, T.-J., and R.-C. Yu, 2005: Atmospheric water vapor transport associated with typical anomalous summer rainfall patterns in China. *J. Geophys. Res.*, **110**, D08104, doi: 10.1029/2004JD005413.

Zhou, T.-J., R. Yu, H. Li, and B. Wang, 2008: Ocean forcing to changes in global monsoon precipitation over the recent half century. *J. Climate*, **21**, 3833-3852, doi: 10.1175/2008 JCLI2067.1.