
- I Main Topics
 - A Equation of a plane
 - B Pole to a plane using cross-products
 - C Intersection of two planes in a line (apparent dip problem)
 - D Intersection of three planes in a point (solution of simultaneous linear equations)

- II Equation for plane: $\mathbf{n} \cdot \mathbf{v} = \mathbf{d}$
 - A Meaning
 - 1 The distance from a known reference point to a plane, as measured in the direction of a unit vector **n** normal to the plane, is d.
 - 2 d > 0 if **n** points from the reference point towards the plane;
 - 3 d < 0 if **n** points from the plane towards the reference point.



- II Equation for plane: $\mathbf{n} \cdot \mathbf{v} = d$ (cont.)
 - B Solution for $\mathbf{n} = \alpha \mathbf{i} + \beta \mathbf{j} + \gamma \mathbf{k} = \langle \alpha, \beta, \gamma \rangle$
 - 1 If x=north, y=east, z=down $\alpha = \cos \phi \cos \theta$, $\beta = \cos \phi \sin \theta$, $\gamma = \sin \phi$
 - 2 If x=east, y=north, z=up $\alpha = cos \varphi sin \theta, \ \beta = cos \varphi cos \theta, \ \gamma = -sin \varphi$ where θ is the pole trend and φ is the pole plunge
 - 3 Matlab:[alpha,beta,gamma]=sph2cart(trend,plunge,1) Trend and plunge must be in radians.
 - 4 Matlab: [trend,plunge,R]=cart2sph(alpha,beta,gamma)
 - 5 Good for z = down; trend and plunge must be in radians.

II Equation for plane: $\mathbf{n} \cdot \mathbf{v} = d$ (cont.)


$$C \mathbf{v} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k} = \langle x, y, z \rangle$$

D Solution for d


1
$$d = \alpha x + \beta y + \gamma z$$


(Normal form)

2 d = dot(n,v)



- III Pole to a plane using cross-products
 - A Consider three noncolinear points A,B,C that form a <u>clockwise</u> circuit as viewed from above
 - B (**B-A**) x (**C-A**) is normal to the plane
 - C $n = (B-A) \times (C-A)$ $|(B-A) \times (C-A)|$, unit normal to the plane.






IV Intersection of two planes in a line

- A Two planes A and B intersect in a line. The equations of those two planes define the line.
- B The direction of intersection is along the vector **a x b**, where **a** is normal to plane A, and **b** is normal to plane B

Direction of Intersection of Two Planes




a is unit normal to Plane A and lies in Plane Cb is unit normal to Plane B and lies in Plane Cc is unit normal to Plane C

c= axb/laxbl

IV Intersection of two planes in a line (cont.)

- C Apparent dip
 - 1 How a geologic plane appears to dip in a vertical cross section of <u>arbitrary</u> strike
 - 2 Plunge of the line of intersection between a geologic plane and a vertical cross section plane of <u>arbitrary</u> strike
 - 3 Plunge of the cross product of two vectors
 - a Vector normal to a geologic plane
 - b Vector normal to vertical cross section plane of <u>arbitrary</u> strike
 - c Make sure cross product points down to get a pole



Vertical plane that is perpendicular to strike of dipping plane

Apparent Dip

- 1) a/b = tan(true dip).
- 2) $a = b \tan(true dip) = c \tan(app. dip)$. So
- 3) b tan(true dip) = c tan(app. dip)


Now consider the angle Ψ between the line of strike of the cross section and the line of strike for the dipping plane.

- 4) $b/c = \sin \Psi$, so
- $c = b/\sin Ψ$.

Now insert (5) into the right side of (3)

- 6) b tan(true dip) = (b/sinΨ) tan(app. dip)
- 7) app. dip = $tan^{-1}\{[tan(true dip)][sin\Psi]\}$ Since $sin\Psi \le 1$, app. dip $\le true dip$.

If $\Psi = 90^{\circ}$, then app. dip = true dip.





Vertical plane that is perpendicular to strike of dipping plane

IV Intersection of two planes in a line (cont.)

D Apparent dip and orthographic projection example

Find the apparent dip of plane ABC in a vertical cross section that strikes N45W.







- C Apparent dip and orthographic projection example
- (1) In the top view, intersect the cross section plane with plane ABC. Call the line of intersection DE. Point D is on line AC. Point E is on line BC.

Next we find the plunge of line DE.

(2) Project the line of intersection DE onto an auxiliary view that is taken parallel to the line of cross section. The plunge of line DE is the apparent dip of plane ABC as seen in the cross section.






V Intersection of features in a point

- A Intersection of two lines (L₁ and L₂) at a point 1 Intersection criteria
 - a Vectors along the lines are not parallel (i.e., $\vec{L}_1 \times \vec{L}_2 \neq 0$)
 - b Vectors along the lines (and normals to the lines) lie in the same plane:

$$\left(\vec{n}_1 \times \vec{L}_1\right) \times \left(\vec{n}_2 \times \vec{L}_2\right) = 0$$




A Intersection of two lines (cont.)

2 Write equations for two lines (n•v=d)

$$\vec{n}_1 \bullet \vec{v}_1 = a_{n_1 x} x + a_{n_1 y} y = d_1$$

$$\vec{n}_2 \bullet \vec{v}_2 = a_{n_2 x} x + a_{n_2 y} y = d_2$$

3 Find n₁, n₂, d₁, d₂



Example

$$(0)(-1)+(1)(2) = 2 = d_1$$


(1)(1.5)+(0)(-1)=1.5 = d₂

A Intersection of two lines (cont.)

3 Write equations for lines in matrix form

$$\begin{bmatrix} a_{n_1x} & a_{n_1y} \\ a_{n_2x} & a_{n_2y} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} d_1 \\ d_2 \end{bmatrix}$$
$$\begin{bmatrix} A \end{bmatrix} \quad \begin{bmatrix} X \end{bmatrix} = \begin{bmatrix} B \end{bmatrix}$$

We seek the vector [X] that satisfies both equations (i.e., a point (x,y) on both lines)



Example

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 1.5 \end{bmatrix}$$
$$[A] \quad [X] = [B]$$

A Intersection of two lines (cont.)

4 Solve for [X]


$$[A][X] = [B]$$

$$[A]^{-1}[A][X] = [A]^{-1}[B]$$

$$[X] = [A]^{-1}[B]$$

1.5000

a Matlab



b Cramer's Rule


$$x = \frac{\begin{vmatrix} B_1 & A_{12} \\ B_2 & A_{22} \end{vmatrix}}{\begin{vmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{vmatrix}} = \frac{\begin{vmatrix} 1.5 & 1 \\ 2 & 0 \end{vmatrix}}{\begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix}} = \frac{-2}{-1} = 2$$

$$y = \frac{\begin{vmatrix} A_{11} & B_1 \\ A_{21} & B_2 \end{vmatrix}}{\begin{vmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{vmatrix}} = \frac{\begin{vmatrix} 0 & 1.5 \\ 1 & 2 \end{vmatrix}}{\begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix}} = \frac{-1.5}{-1} = 1.5$$

2.0000

- B Intersection of three plane (P₁, P₂, and P₃) at a point
 - 1 Intersection criteria
 Normals of any planes are not parallel; scalar triple product of normals to planes ≠ 0

(i.e., $n_1 \bullet (\vec{n}_2 \times \vec{n}_3) \neq 0$)



B Intersection of three planes (cont.)

2 Write equations for three planes

$$(n \cdot v = d)$$

$$\vec{n}_1 \bullet \vec{v}_1 = a_{n_1 x} x + a_{n_1 y} y + a_{n_1 z} z = d_1$$

$$\vec{n}_2 \bullet \vec{v}_2 = a_{n_2 x} x + a_{n_2 y} y + a_{n_2 z} z = d_2$$

$$\vec{n}_3 \bullet \vec{v}_3 = a_{n_3 x} x + a_{n_3 y} y + a_{n_3 z} z = d_3$$


3 Find n_1 , n_2 , n_3 , d_1 , d_2 , d_3

Example

$$n_1 = 1\vec{i} + 0\vec{j} + 0\vec{k}$$

$$n_2 = 0\vec{i} + 1\vec{j} + 0\vec{k}$$

$$n_3 = 0\vec{i} + 0\vec{j} + 1\vec{k}$$



$$(1)(3)+(0)(y)+(0)(z)=3=d_1$$

$$(0)(x)+(1)(2)+(0)(z)=2=d_2$$

$$(0)(x)+(0)(y)+(1)(1)=1=d_2$$


B Intersection of three planes (cont.)

3 Write equations for lines in matrix form

$$\begin{bmatrix} a_{n_{1}x} & a_{n_{1}y} & a_{n_{1}z} \\ a_{n_{2}x} & a_{n_{2}y} & a_{n_{2}z} \\ a_{n_{3}x} & a_{n_{3}y} & a_{n_{3}z} \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} d_{1} \\ d_{2} \\ d_{3} \end{bmatrix}$$

$$[A] \qquad [X] = [B]$$

planes)



We seek the vector [X] that satisfies each equation (i.e., one point (x,y,z) on all three planes)
$$\begin{bmatrix}
1 & 0 & 0 \\
0 & 1 & 0 \\
0 & 0 & 1
\end{bmatrix}
\begin{bmatrix}
x \\
y \\
z
\end{bmatrix} = \begin{bmatrix}
3 \\
2 \\
1
\end{bmatrix}$$

$$[A] \quad [X] = [B]$$

B Intersection of three planes (cont.)

4 Solve for [X]


$$[A][X] = [B]$$

$$[A]^{-1}[A][X] = [A]^{-1}[B]$$

$$[X] = [A]^{-1}[B]$$


a Matlab

9/20/13



b Cramer's Rule
Could use Cramer's
rule, but it is
cumbersome.

- C Intersection of a line and a plane
 - 1 A line (L₁) and a plane (P₁) intersect at a point
 - 2 Point of intersection can be viewed as the intersection of 3 planes
 - a Plane P₁ (white)
 - Plane P₂ (blue)
 P₂ intersects plane P₃ at line
 L₁. Plane P₂ is a vertical plane containing L₁, so P₂ strikes parallel to the trend of L₁.
 - c Plane P_3 (gray) P_3 intersects plane P_2 at line P_3 . Plane P_3 is an inclined plane that contains P_3 . The dip of P_3 equals the plunge of P_3 , and the strike of P_3 is 90° from the trend of P_3 .



Two Math Handbook References

Gellert, W., Küstner, H., Hellwich, M,. and Kästner, H., 1977, The VNR concise encyclopedia of mathematics: Van Nostrand Reinhold, New York, 760 p.

Tuma, J.J., 1979, Engineering mathematics handbook: McGraw-Hill, New ork, 394 p.