

FOR IMMEDIATE RELEASE

SOEST Media Contact: Tara Hicks Johnson
808-956-3151, hickst@hawaii.edu

Ocean's 'Twilight Zone' Plays Important Role in Climate Change

New study identifies a critical link influencing the ocean's ability to store carbon dioxide


Honolulu, HI - A major study has shed new light on the dim layer of the ocean called the "twilight zone"—where mysterious processes affect the ocean's ability to absorb and store carbon dioxide accumulating in our atmosphere.

The results of two international research expeditions to the Pacific Ocean, published April 27 in the journal *Science*, show that carbon dioxide—taken up by photosynthesizing marine plants in the sunlit ocean surface layer—does not necessarily sink to the depths, where it is stored and prevented from re-entering the atmosphere as a greenhouse gas. Instead, carbon transported to the depths in sinking particles is often consumed by animals and bacteria and recycled in the twilight zone—100 to 1,000 meters below the surface—and never reaches the deep ocean.

Using new technology, the researchers found that only 20 percent of the total carbon in the ocean surface made it through the twilight zone off Hawaii, while 50 percent did in the northwest Pacific near Japan.

The twilight zone acts as a "gate," allowing more sinking particles through in some regions and fewer in others, complicating scientists' ability to predict the ocean's role in offsetting the impacts of greenhouse gases. It also adds a new wrinkle to proposals to mitigate climate change by fertilizing the oceans with iron—to promote blooms of photosynthetic marine plants and transfer more carbon dioxide from the air to the deep ocean.

"The twilight zone is a critical link between the surface and the deep ocean," said Ken Buesseler, a biogeochemist at Woods Hole Oceanographic Institution (WHOI) and lead author of the new study in *Science*, co-authored by 17 other scientists, including two from the University of Hawaii. "We're interested in what happens in the twilight zone, what sinks into it and what actually sinks out of it. Unless the carbon that gets into the ocean goes all the way down into the deep ocean and is stored there, the carbon can still make its way back into the atmosphere. Without long-term carbon storage at depth, the ocean can do little to stem the increase in atmospheric levels of carbon dioxide, a greenhouse gas that impacts the earth's climate."



Particles sinking from sunlit surface waters through the ocean's dimly lit twilight zone are swept sideways by currents. Conventional moored or tethered traps designed to catch the particles are like "rain gauges in hurricanes," said WHOI biogeochemist Ken Buesseler. He and engineer Jim Valdes are designing a new-generation neutrally buoyant untethered vehicle called the Twilight Zone Explorer, which will be swept along with the currents. It will surface periodically to relay data via satellite. (Illustration by Jack Cook, Woods Hole Oceanographic Institution)



Neutrally Buoyant Sediment Trap being lowered into water from RV Kilo Moana off Hawaii. (Photo by Ken Buesseler, Woods Hole Oceanographic Institution)

Buesseler was a leader of the ambitious project, funded primarily by the US National Science Foundation (NSF), called VERTIGO (VERTical Transport In the Global Ocean). More than 40 biologists, chemists, physical oceanographers, and engineers from 14 institutions and seven countries participated in the two VERTIGO cruises in 2004 and 2005 to investigate how marine plants die and sink, or are eaten by animals and converted into sinking fecal pellets.

These sinking particles, often called “marine snow,” supply food to organisms deeper down, including bacteria that decompose the particles. In the process, carbon is converted back into dissolved organic and inorganic forms that are re-circulated and reused in the twilight zone and that can make their way to the surface and back into the atmosphere.

One of the sampling sites was Station ALOHA, the NSF-funded Hawaii Ocean Time-series (HOT) research site. Since 1988, HOT investigators have catalogued information about the biologic, chemical and physical parameters of the surrounding ocean, making it one of the most comprehensively characterized sites on our planet.

Both the VERTIGO and HOT programs had several noteworthy roots, none as important as the VERTical Transport and EXchange (VERTEX) program that ran from about 1980-1988, just as HOT was getting started. VERTEX pioneered many new methods of analysis and provided novel data sets on the chemistry and microbiology of the 'twilight zone'. Dave Karl, one of the researchers in this study, was one of the principal investigators in VERTEX which conducted field experiments in the NE Pacific including several stations off Hawaii. “The HOT program established station ALOHA in October 1988 and continued many of the investigations that began during VERTEX, including but not limited to the export of particulate matter from the surface layers of the ocean and the relationship between primary production and export,” says Karl. “These observations from station ALOHA/HOT were used, in part, to develop the hypotheses that were tested during the VERTIGO program. Therefore in many ways VERTIGO was built on a HOT foundation.”

The sites off Hawaii and Japan were selected because they had been the focus of long-term ocean observations by co-authors David Karl (University of Hawaii) and Makio Honda (Japan Agency for Marine-Earth Science and Technology). Oceanographer Robert Bidigare (University of Hawaii) made measurements of phytoplankton pigments on samples collected during both cruises. Biologists Deborah Steinberg (Virginia Institute of Marine Sciences) and Mary Silver (University of California, Santa Cruz) were needed to identify plankton species and understand differences in the food webs that propel the marine carbon cycle. Thomas Trull (University of Tasmania, Australia), Philip Boyd (University of Otago, New Zealand), and Frank Dehairs (Free University of Brussels, Belgium) all study the Southern Ocean and could provide important perspectives contrasting ocean carbon cycle off Antarctica and in VERTIGO. David Siegel (University of California, Santa Barbara) helped track the ocean currents and pathways of the sinking particles. James Bishop (Lawrence Berkeley National Laboratory and University of California, Berkeley) was funded by the US Department of Energy to deploy new autonomous optical sediment traps designed to follow the hourly changes in sedimentation as well as ship deployed particle sampling systems to quantify the abundance and composition of particles in the twilight zone.

“This combination of expertise could not be found in any single lab or country,” Buesseler said. “We were fortunate to attract such a diverse group of talented scientists willing to unravel the secrets of the twilight zone and its role in the global carbon cycle.”

While many studies have investigated the surface of the ocean, little research has been conducted on the carbon cycle below. The VERTIGO team examined a variety of processes to open a new window into the difficult-to-explore twilight zone. They successfully used a wide array of new tools, including an experimental device that overcame a longstanding problem of how to collect marine snow falling into the twilight zone.


The problem is that particles sink slowly, perhaps 10 to a few hundred meters per day, but they are swept sideways by ocean currents traveling many thousands of meters per day. To collect sinking particles, scientists use cones or tubes that hang beneath buoys or float up from seafloor. That, Buesseler said, “is like putting out a rain gauge in a hurricane.”

Buesseler and WHOI engineer Jim Valdes developed Neutrally Buoyant Sediment Traps (NBST)—free-floating devices that sink to a programmed depth within the twilight zone and neither sink nor rise. They are swept along with the currents for several days, collecting particles, and then programmed to resurface, transmit their position via satellite, and wait for recovery, more than 10 to 20 miles away from where they were dropped into the ocean.

“It’s a bit like finding a needle in a haystack, since they are so small and difficult to spot, especially in rough seas that are common in the open ocean,” Valdes said.

Water column and sediment trap samples collected on both cruises were returned to Robert Bidigare’s lab at the University of Hawaii for analysis. The goal was to determine the composition of phytoplankton (the “primary producers”) at both study sites as well as the amount and makeup of phytoplankton removed from the upper ocean by sinking and zooplankton grazing processes.

On their first scientific mission for VERTIGO in 2004, Buesseler and Valdes could only wait and hope their devices would work. “Seven NBSTs went in the water and all seven came back with their precious cargo—a first in ocean sciences history,” Buesseler said.



A humorous plug for VERTIGO was incorporated into the analysis of zooplankton fecal pellets collected from sediment traps at ALOHA and K2. (Image by Stephanie E. Wilson, VIMS)

Why more carbon reached the depths in the northwest Pacific might be due to many factors. Waters there are full of silica that plankton incorporate to make shells. Do the silica-laden plankton weigh more and thus sink faster, giving bacteria less time to break them down? Do lower water temperatures in the northwest Pacific slow down the breakdown of organic carbon? Do different populations in the food webs at different sites change how organic matter is broken down and marine snow is produced? Scientists will continue to explore these regional differences in the ability of carbon to reach the deep sea.

The next step for the Hawaii researchers will be a major research cruise on the University of Hawaii’s research vessel the R/V Kilo Moana. “As we learn more and more about the sea around us, including the deep ocean, we gain an appreciation for how important and how fragile life in these regions can be,” says Karl. “By gaining a better understanding of how organic matter and energy are processed in the mid-depths of the ocean we can better predict how life in the sea may be affected by human-imposed climate change including greenhouse gas production. The twilight zone may be the largest habitat for life on our planet in terms of volume, yet we are just beginning to understand who lives there and how they make their living.”

WHOI Media Release, with additional information from the SOEST Outreach Office.

Images available, please contact Tara Hicks Johnson, Outreach Specialist, School of Ocean & Earth Sciences & Technology, Phone (808) 956-3151, email hickst@hawaii.edu

For Interviews contact:

Robert R. Bidigare, Department of Oceanography, School of Ocean & Earth Science and Technology, University of Hawaii at Manoa, Honolulu, HI Phone: (808) 956-8146, bidigare@hawaii.edu

David Karl, Department of Oceanography, School of Ocean & Earth Science and Technology, University of Hawaii at Manoa, Honolulu, HI Phone: (808) 956-8964, dkarl@hawaii.edu

Related University of Hawaii links:

Hawaii Ocean Time-series http://hahana.soest.hawaii.edu/hot/hot_jgofs.html

Center for Microbial Oceanography: Research and Education (C-MORE) <http://cmore.soest.hawaii.edu/>

Related WHOI links:

WHOI Contact: Media Relations Office, 508-289-3340, media@whoi.edu

A Journey to the Ocean’s Twilight Zone – Oceanus Magazine www.whoi.edu/oceanus/viewArticle.do?id=15328

Swimming in the Rain – Oceanus Magazine www.whoi.edu/oceanus/viewArticle.do?id=15329

VERTIGO - Ocean Particle Fluxes www.whoi.edu/science/MCG/cafethorium/website/projects/vertigo.html